


HOTCHKISS (L.W.)

Report of a Case in which
LAPAROTOMY
was performed for Intestinal Perfor-
ation occurring in the Course
of Typhoid Fever.

BY

L. W. HOTCHKISS, M. D.,
SURGEON TO THE MANHATTAN HOSPITAL, ETC.

REPRINTED FROM THE
New York Medical Journal
for January 11, 1896.



*Reprinted from the New York Medical Journal
for January 11, 1896.*



REPORT OF A CASE IN WHICH
LAPAROTOMY
WAS PERFORMED FOR INTESTINAL PERFORATION
OCCURRING IN THE COURSE OF
TYPHOID FEVER.*

By L. W. HOTCHKISS, M. D.,
SURGEON TO THE MANHATTAN HOSPITAL, ETC.

ON the 13th of August, 1895, I was asked to see with Dr. Knickerbocker, at the Manhattan Hospital, a case in which he had made the diagnosis of intestinal perforation. The patient was a Swede, twenty-four years of age, a carpenter by trade, who had been admitted to the hospital, August 7th, with the following history: He had been in the hospital a year before, suffering from an intermittent fever, for which he had been treated and discharged cured. His present illness began about ten days before his admission, with chills, fever, headache, and anorexia, accompanied with great prostration. He had not been confined to bed all the time, however, and, though he felt very wretchedly, had managed to walk to the hospital. The examination of heart and lungs was negative. There were several small, rose-colored spots over the chest and abdomen, which disappeared on pressure. The

* Read before the Society of Alumni of Bellevue Hospital, November 6, 1895.

temperature was 103.8° F. A diagnosis of typhoid fever was made, and treatment by Brandt's method of baths instituted. The patient bore the baths well, his tongue grew moist, and his pulse improved.

On the night of August 12th he complained suddenly of intense abdominal pain, and was ordered morphine. Next morning he was seen by Dr. Knickerbocker, the attending physician, who suspected perforation, and asked me to see the case with him, with a view to operation if that should be deemed advisable.

On examination, the abdomen was found considerably distended, exquisitely tender on pressure, and the patient suffering constantly from general abdominal pain. His face was anxious and drawn; his intelligence perfectly good. He had vomited. Pulse rapid; temperature high. The seriousness of his condition was presented to him, and the possibility of relief by surgical means explained to him. He said he was anxious to have something done, and would willingly submit to operation to obtain any relief from his terrible pain. He was transferred to the surgical side, and operated upon as soon as preparations could be made.

Ether was used as an anæsthetic, and a median incision, three inches and a half in length, made between the umbilicus and the pubes. On incising the peritonæum, a large quantity of cloudy, serous fluid, containing big flakes of fibrin and probably some fecal matter, gushed out. There was but little odor to this fluid. The peritonæum covering the small intestines and everywhere was deeply congested, and covered here and there by patches of soft fibrin. There were no adhesions. Enlarged lymphatic glands in the mesentery were distinctly visible and palpable. The cavity was flushed out thoroughly with hot normal salt solution, and a search for the perforation was at once made. It was easily found in the lower part of the ileum, about five inches above the ileo-cæcal junction and on the anterior wall of the gut. The perforation was quite small, its edges ragged, and the tissues surrounding it infiltrated and much softened. Through the wall of the gut the thickening, corresponding apparently to the enlarged

Peyer's patch, could be felt. The perforation and softened area about were turned in longitudinally for an inch or more and sutured with fine black silk after Lembert's method. The turning in of the intestinal wall well beyond the edges of the ulcer brought into contact a considerable area of peritoneal surface. After again thoroughly flushing the whole peritoneal cavity with hot sterile normal salt solution, the diseased portion of the ileum was surrounded and isolated by packing of sterile gauze. The upper end of the abdominal wound was closed, the lower left open for the gauze drainage. After the operation the pulse continued very weak, but the patient recovered consciousness and said his pain was less. He continued to sink, however, and died about four hours and a half after the operation, his temperature rising to 108° F.

An examination of the abdominal cavity after death showed essentially the same conditions as described. There were no other perforations found, and there had apparently been no leakage from the sutured rent.

The portion of the ileum which included the sutured area was removed and sent to the Carnegie laboratory. Sections through the sutured portion of gut showed that sufficient fibrin had already been thrown out to hold the sutured peritoneal surfaces in firm apposition.

In putting on record this case in which laparotomy was done for perforating typhoid ulcer, one more is added to the small number of cases in which an attempt has been made to relieve this fatal complication of typhoid fever by surgical means.

According to a recent article by Dr. F. H. Wiggin, there have been only seventeen well-authenticated cases of perforation of the gut in typhoid fever reported in which a laparotomy was performed, and out of these seventeen patients so operated upon three recovered.

Of the three successful cases, one belongs to Dr. Van Hook, a second to a Russian surgeon, Dr. Netschajan, the third to Dr. Robert Abbe, of this city.

4 LAPAROTOMY FOR INTESTINAL PERFORATION.

Realizing that in most of these cases we have to deal with a very rapid and fatal form of peritonitis, our only hope seems to lie in an early operation. In the case reported, about ten or twelve hours had elapsed since the perforation had occurred, and general septic peritonitis had supervened, rendering the case practically hopeless. I feel that in perforations of the gut in typhoid fever we may not always have to deal with an equally rapid and fatal infection. As in perforative appendicitis different forms of peritonitis may result, varying from a fibro-purulent peritonitis more or less extensive to an acute septic general peritonitis or peritoneal sepsis which is rapidly fatal, so in typhoid fever it would seem that in some cases where the ulcerative process is not too rapid and the infection not too intense and overwhelming, a peritonitis might result which could be as successfully treated by an early laparotomy as are many cases of this disease due to perforative appendicitis.

Realizing that in these cases we have to work with the chances of a successful issue decidedly against us, and upon a patient in the worst possible condition generally for operation, I feel that it is only fair to give the patient the possible chance, small though it be. I feel that the operation is not only justifiable in many cases, but should be offered to the patient as giving some hope in a condition otherwise practically without it.

I think the operation should be classed with tracheotomy for laryngeal stenosis, and with herniotomy for the relief of strangulation—*i. e.*, as an occasional life-saving procedure. It is not too much to expect, perhaps, that some time in the near future the surgeon will be oftener called in cases of typhoid fever where the diagnosis of perforation is made.

The New York Medical Journal.

A WEEKLY REVIEW OF MEDICINE.

EDITED BY

FRANK P. FOSTER, M.D.

THE PHYSICIAN who would keep abreast with the advances in medical science must read a *live* weekly medical journal, in which scientific facts are presented in a clear manner; one for which the articles are written by men of learning, and by those who are good and accurate observers; a journal that is stripped of every feature irrelevant to medical science, and gives evidence of being carefully and conscientiously edited; one that bears upon every page the stamp of desire to elevate the standard of the profession of medicine. Such a journal fulfills its mission—that of educator—to the highest degree, for not only does it inform its readers of all that is new in theory and practice, but, by means of its correct editing, instructs them in the very important yet much-neglected art of expressing their thoughts and ideas in a clear and correct manner. Too much stress can not be laid upon this feature, so utterly ignored by the “average” medical periodical.

Without making invidious comparisons, it can be truthfully stated that no medical journal in this country occupies the place, in these particulars, that is held by THE NEW YORK MEDICAL JOURNAL. No other journal is edited with the care that is bestowed on this; none contains articles of such high scientific value, coming as they do from the pens of the brightest and most learned medical men of America. A glance at the list of contributors to any volume, or an examination of any issue of the JOURNAL, will attest the truth of these statements. It is a journal for the masses of the profession, for the country as well as for the city practitioner; it covers the entire range of medicine and surgery. A very important feature of the JOURNAL is the number and character of its illustrations, which are unequaled by those of any other journal in the world. They appear in frequent issues, whenever called for by the article which they accompany, and no expense is spared to make them of superior excellence.

Subscription price, \$5.00 per annum. Volumes begin in January and July.

PUBLISHED BY

D. APPLETON & CO., 72 Fifth Avenue, New York.

